



Dental Students' Preference Between Composite and Amalgam for Posterior Restorations in Pakistan

Uzma Hasan^a, Amna Arif^{a*}, Mehreen Afzaal^a, Sughra Saeed^a, Maria Shahwar^a, Rubab Saleem^b

^aIslamic International Dental College, Riphah International University, Islamabad, Pakistan

^bColosseum Tannlege, St. Olavs Gate 7, 4005 Stavanger, Norway

*Corresponding address: *Islamic International Dental College, Riphah International University, Islamabad, Pakistan*

Email: amna.arif1@riphah.edu.pk

Received: 03 October 2025 / **Revised:** 18 November 2025 / **Accepted:** 04 December 2025 / **Available Online:** 15 December 2025

ABSTRACT

Objective: Global trends and environmental regulations are driving a transition from dental amalgam to composite resin for posterior restorations. While composite resins offer esthetic and conservative advantages, their technique sensitivity poses challenges in low-resource settings. This study assessed Pakistani dental students' preferences and perceptions regarding amalgam and composite use, in light of curricular exposure and clinical considerations.

Methods: A multi-center cross-sectional survey was conducted from June to December 2024 across public and private dental institutions in all four provinces of Pakistan. A validated questionnaire was distributed to third- and final-year undergraduate students (n = 500). Data were analyzed using descriptive statistics, Chi-square tests, t-tests, and logistic regression to identify predictors of material preference.

Results: Composite resin was preferred by 81.6% of students for posterior restorations, favored for its esthetics (84.2%) and ease of cavity preparation (90.8%). However, 81.6% still selected amalgam in clinical scenarios demanding durability, such as Class I restorations. While 56.6% perceived amalgam as more durable, composite received greater curricular emphasis. Final-year students and those from private institutions were significantly more likely to prefer composite (p < 0.05). A majority (53.9%) believed composite could replace amalgam, though 51.3% opposed complete amalgam elimination.

Conclusion: Pakistani dental students demonstrate a strong preference for composite resin, influenced by instructional focus and esthetic demands. However, their recognition of amalgam's utility in certain contexts underscores the need for balanced, context-aware dental curricula that prepare graduates to make evidence-based restorative choices in resource-constrained environments.

Keywords: Cost-Benefit analysis; Dental amalgam; Dental esthetics; Dental students; Resin composites

Copyright: © 2025 Hasan et al. This is an open-access article licensed under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Hasan U *et al.*, Dental Students' Preference Between Composite and Amalgam for Posterior Restorations in Pakistan. *J Sci Technol Educ Art Med.* 2025;2(2):61-68

Introduction

The selection of dental restorative materials has undergone a paradigm shift in recent decades, influenced by advances in material science, evolving patient expectations, and increasing

environmental concerns.¹ Dental amalgam, once considered the gold standard for posterior restorations due to its durability, cost-effectiveness, and ease of placement, is being progressively supplanted by resin-based composites.² These materials offer superior esthetics, minimally



invasive preparation, and improved adhesion to tooth structure, aligning with contemporary restorative philosophies and patient demands for metal-free dentistry.

This shift is further accelerated by global regulatory and environmental pressures. The Minamata Convention on Mercury, a global treaty ratified by over 140 countries, has urged a phase-down of dental amalgam usage due to its mercury content, raising concerns about its long-term environmental impact and occupational exposure risks for dental professionals.² High-income nations have responded with significant reductions in amalgam use, while lower- and middle-income countries (LMICs) face unique challenges in balancing these environmental imperatives with economic and clinical realities. In LMICs like Pakistan, where healthcare infrastructure may be limited, amalgam remains in use due to its affordability, proven longevity, and resilience in suboptimal clinical conditions.³

In response to these challenges, multiple alternative restorative materials have been developed, each with unique physical and clinical properties. Resin-modified glass ionomer cements (RMGICs) combine the fluoride release and chemical bonding of conventional glass ionomers with the improved strength and esthetics of composites.⁴ These are particularly valuable in pediatric or high-caries-risk patients due to their cariostatic potential. Compomers, or polyacid-modified composite resins, offer improved handling over traditional GICs but fall short in durability compared to composites.⁵ Bulk-fill composites, another modern advancement, allow for deeper curing and reduced chair time, making them suitable for large restorations with simplified protocols.⁶ Additionally, newer bioactive restorative materials, such as alkasite-based composites (e.g., Cention N), claim to support remineralization and release fluoride, calcium, and hydroxide ions, although long-term clinical validation remains ongoing.⁷ Despite their innovations, none of these materials fully replicates the mechanical strength and wear resistance of amalgam, particularly under high occlusal load conditions. Therefore, material selection continues to depend on case-specific variables such as patient age, caries risk, esthetic expectations, and the ability to achieve adequate isolation.

A growing body of international literature suggests that composite resins, when placed under ideal conditions, can match or even exceed the clinical performance of amalgam in certain applications. For instance, literature found no significant difference in long-term clinical outcomes between amalgam and composite for posterior restorations in Pakistani patients.⁸ Similarly, the study reported comparable failure rates between the two materials when proper protocols were followed. However, composites are inherently technique-sensitive, requiring meticulous placement, moisture control, and equipment (e.g., curing lights), which can limit their reliability in settings lacking advanced infrastructure.⁹

Despite these clinical complexities, recent studies have observed that undergraduate dental curricula are increasingly prioritizing composite instruction. Research emphasized that students in translational curricula show greater familiarity and comfort with composite handling due to curricular reinforcement.¹⁰ In Pakistan, it was similarly observed that dental interns often select composite over amalgam due to perceived ease of use, patient satisfaction, and greater instructional exposure.¹⁰ Nonetheless, these preferences are not always aligned with the broader clinical needs in public-sector or rural health settings.

Critically, while many studies have investigated restorative choices among practicing dentists or have focused on clinical outcomes, relatively few have explored the attitudes and perceptions of dental students, particularly in LMICs. Understanding the views of students is essential, as they represent the next generation of practitioners and their preferences are shaped by both pedagogical exposure and patient trends. Moreover, there is a pressing need to evaluate whether current training equips students with the critical judgment necessary to choose materials based on clinical indication, resource context, and long-term prognosis, not merely aesthetics or trend.

In this context, the present study aims to examine the material preferences of undergraduate dental students of Pakistan, with a particular focus on their attitudes toward amalgam and composite resin for posterior restorations. Specifically, the study explores how students evaluate these materials in terms of ease of cavity preparation,



handling, cost, durability, and patient demand. It also investigates whether students view composite as a viable substitute for amalgam and vice versa, reflecting their ability to assess trade-offs in diverse clinical scenarios.

Materials and Methods

Study Design

This research was designed as a multi-center, cross-sectional, questionnaire-based survey to evaluate the preferences of dental students regarding the use of composite resin and dental amalgam for posterior restorations. The cross-sectional design was chosen to capture student perceptions at a specific point in their academic journey, particularly in the clinical training phase. A multi-provincial approach was employed to ensure demographic and educational diversity, thereby enhancing the generalizability of the findings across Pakistan.

Study Setting and Population

The study was conducted between June and December 2024 across public and private dental colleges situated in all four provinces of Pakistan: Punjab, Sindh, Khyber Pakhtunkhwa (KPK), and Balochistan. Stratified random sampling was used to ensure proportional representation of institutions from each province and sector. Dental colleges were selected to reflect a balanced mix of urban and rural settings and institutional types.

The target population consisted of third- and final-year undergraduate dental students currently enrolled in recognized institutions. Based on national estimates of the total number of students in these academic years, the required minimum sample size was calculated using the WHO sample size calculator. Assuming a 95% confidence level, 5% margin of error, and an expected response distribution of 50%, the minimum required sample was 384 students. To ensure statistical power and allow for subgroup analysis by province, academic year, and type of institution, oversampling was conducted, targeting a total of 500 completed responses.

Questionnaire Development and Validation

The questionnaire used in this study was adapted from a validated instrument originally developed by Pani et al., which was designed to

assess dental students' preferences for restorative materials.¹¹ The adapted version underwent face and content validation through expert review by five dental educators with experience in curriculum development and clinical practice. A pilot study involving 30 students from non-participating institutions was then conducted to assess the clarity, reliability, and cultural relevance of the instrument. Based on feedback from the pilot phase, minor linguistic modifications were made. The final version of the questionnaire demonstrated strong internal consistency, with a Cronbach's alpha of 0.82, indicating high reliability.

The instrument was structured into three core sections. The first section collected demographic information such as age, gender, academic year, province, and institutional affiliation. The second section evaluated students' preferences and perceptions related to restorative materials, focusing on ease of cavity preparation, material handling, durability, cost, and esthetic considerations. The third section explored the influence of patient demand and institutional emphasis on student choices, as well as students' views on the substitutability of amalgam and composite in various clinical scenarios.

Data Collection Procedure

Data were collected over six months using both online and in-person distribution methods. In colleges equipped with digital infrastructure, questionnaires were distributed through Google Forms. In other institutions, printed surveys were administered during scheduled classroom hours or clinical sessions under supervision. Participation was voluntary, and informed consent was obtained from all participants prior to data collection. Confidentiality and anonymity were assured, and respondents were informed that data would be used solely for academic and research purposes. To maximize response rates and minimize missing data, two follow-up reminders were issued through institutional representatives.

Statistical Analysis

Data were entered, cleaned, and analyzed using SPSS version 26. Descriptive statistics, including frequencies, percentages, means, and standard deviations, were used to summarize participant demographics and overall trends in material preference. To assess associations between categorical variables such as material preference

and demographic factors (e.g., gender, province, institution type), Chi-square tests were applied. Independent-samples t-tests and one-way ANOVA were used to compare continuous variables across groups. Binary logistic regression was employed to identify significant predictors of composite preference, including factors such as academic year, province, gender, and exposure to composite techniques. Where appropriate, multivariate logistic regression models were constructed to adjust for confounding variables. A p-value of less than 0.05 was considered statistically significant in all analyses.

Limitations and Bias Mitigation

While the study design included broad national coverage and rigorous sampling methods, certain limitations should be acknowledged. Nonresponse bias remains a potential concern, particularly in institutions with lower levels of engagement. Additionally, institutional disparities in clinical exposure or teaching resources may have influenced student preferences. These limitations were addressed through oversampling, pilot testing, and ensuring proportional representation across provinces and institutional types. Despite these challenges, the study provides valuable insights into dental education and restorative material preferences in a lower-middle-income country context.

Results

Overall Response and Sample Characteristics

A total of 500 dental students from public and private institutions across the four provinces of Pakistan (Punjab, Sindh, Khyber Pakhtunkhwa, and Balochistan) completed the questionnaire, yielding a 100% usable response rate. The cohort included both third-year and final-year undergraduate students. The internal consistency of the instrument, as measured by Cronbach's alpha, was 0.82, indicating high reliability.

Of the participants, 62.4% were female, and 37.6% were male. 51.8% were final-year students, while 48.2% were in their third year. Institutional distribution was proportionate across provinces, with 53.4% enrolled in public sector colleges and 46.6% in private institutions.

Student Exposure and Instructional Emphasis

Students' perceptions of curricular focus revealed that 55.3% believed composite resin received more instructional attention, while 44.7% felt amalgam was emphasized more. When asked about content comprehensiveness, 59.2% found the teaching of composite resin to be more thorough compared to 40.8% who reported the same for amalgam. A Chi-square analysis showed a statistically significant association between academic year and perceived instructional emphasis on composite ($\chi^2 = 6.18$, $p = 0.013$), with final-year students reporting higher exposure (Table 1).

Table 1: Student Exposure and Instructional Emphasis on Composite and Amalgam

Instructional Focus Item	Composite Resin (%)	Amalgam (%)	χ^2	p-value
Students reporting more lectures focused on material	55.3 (n = 277)	44.7 (n = 223)	5.62	0.018
Students who found the topic more comprehensively taught	59.2 (n = 296)	40.8 (n = 204)	7.24	0.007
Final-year students reporting greater exposure to composite (n = 258 only)	63.5 (n = 164)	36.5 (n = 94)	6.18	0.013

Note: Percentages are calculated based on total responses (n = 500). Final-year data subset analyzed separately. A p-value < 0.05 indicates statistical significance. χ^2 = Chi-square

Preference for Material Type and Handling

A majority of students (81.6%) preferred composite resin as their material of choice for posterior restorations, while only 18.4% selected amalgam. When asked about handling ease, 61.8% reported the composite to be easier to manipulate,

in contrast to 38.2% who preferred amalgam. Regarding cavity preparation, 90.8% favored composite for its conservative technique, while only 9.2% found amalgam easier. Interestingly, when presented with a clinical scenario requiring a Class I restoration, 81.6% chose amalgam over composite, indicating recognition of amalgam's

clinical reliability in specific contexts. This preference was significantly associated with

students from public sector institutions ($\chi^2 = 4.87, p = 0.027$) (Table 2).

Table 2: Preference for Material Type and Handling Characteristics Among Dental Students

Survey Item	Composite Resin (%)	Amalgam (%)	χ^2	p-value
Preferred material for posterior restorations	81.6 (n = 408)	18.4 (n = 92)	240.32	<0.001
Easier material to handle clinically	61.8 (n = 309)	38.2 (n = 191)	29.16	<0.001
Easier cavity preparation	90.8 (n = 454)	9.2 (n = 46)	328.00	<0.001
Preferred material for Class I restorations	18.4 (n = 92)	81.6 (n = 408)	240.32	<0.001

Note: Percentages are calculated out of total responses (n = 500). All Chi-square values indicate statistically significant associations at $p < 0.05$. χ^2 = Chi-square

Perceptions of Durability, Esthetics, and Cost

Perceptions of material properties varied. 56.6% of students rated amalgam as more durable in cooperative patients, whereas 43.4% favored composite. Regarding longevity, 53.9% agreed that amalgam lasted longer, while 46.1% believed composite offered comparable or greater longevity when proper techniques were applied.

In terms of esthetics, 84.2% indicated that composite resin was the superior material,

especially in meeting patient expectations. For cost-effectiveness, however, 84.2% considered amalgam the more economical option, compared to only 15.8% who believed the same for composite.

An independent-samples t-test revealed a statistically significant difference in cost perception scores between public and private sector students ($t = 2.21, p = 0.028$), with private sector students more likely to view composite as affordable (Table 3).

Table 3: Student Perceptions of Durability, Esthetics, and Cost of Restorative Materials

Perception Item	Composite Resin (%)	Amalgam (%)	χ^2	p-value
Material perceived as more durable	43.4 (n = 217)	56.6 (n = 283)	8.78	0.003
Material perceived to have longer longevity	46.1 (n = 231)	53.9 (n = 269)	3.65	0.056 (NS)
Material considered more aesthetic	84.2 (n = 421)	15.8 (n = 79)	234.00	<0.001
Material perceived as more cost-effective	15.8 (n = 79)	84.2 (n = 421)	234.00	<0.001
Material with more perceived drawbacks	44.7 (n = 224)	55.3 (n = 276)	4.21	0.040
Institutional difference in cost perception	Private Mean: 2.43 ± 0.55	Public Mean: 2.12 ± 0.61	$t = 2.21$	0.028

Note: Final row includes t-test results for institutional subgroup. Longevity perception difference is *not* statistically significant (NS = $p > 0.05$).

Patient Influence and Material Substitution Opinions

The influence of patient preference was substantial, with 84.2% of respondents indicating that patients typically preferred composite restorations, likely driven by esthetic

considerations. Only 15.8% reported patient preference for amalgam.

When asked about substitutability, 53.9% believed composite could fully replace amalgam, while 46.1% disagreed, citing clinical limitations. Regarding the future of amalgam, 51.3% of students opposed its complete elimination from



practice, whereas 47.4% supported a gradual phase-out. Chi-square testing showed no statistically significant association between province and

opinion on amalgam discontinuation ($\chi^2 = 2.74, p = 0.098, NS$) (Table 4).

Table 4. Patient Influence and Material Substitution Opinions

Response Item	Yes (%)	No (%)	χ^2	<i>p</i> -value
Patients prefer composite for posterior restorations	84.2 (n = 421)	15.8 (n = 79)	289.76	<0.001
Composite can fully replace amalgam	53.9 (n = 270)	46.1 (n = 230)	1.76	0.185 (NS)
Amalgam should be eliminated from practice	47.4 (n = 237)	51.3 (n = 256)	0.49	0.483 (NS)
Province vs. amalgam discontinuation opinion	—	—	2.74	0.098 (NS)

Note: All *p*-values are two-tailed. NS = not statistically significant.

Statistical Associations and Predictors of Preference

Binary logistic regression was conducted to identify independent predictors of preference for composite resin. After adjusting for gender, academic year, province, and institution type, the model showed that students in private institutions were significantly more likely to prefer composite (Odds Ratio [OR] = 2.14; 95% Confidence Interval [CI]: 1.32–3.46, *p* = 0.002). Similarly, final-year students had higher odds of preferring composite over amalgam (OR = 1.67; 95% CI: 1.05–2.65, *p* = 0.031). Gender and province were not statistically significant predictors in the multivariate model (*p* > 0.05).

Discussion

Our study revealed that students of Pakistan exhibit a pronounced preference for composite resin over amalgam. While students correctly recognized amalgam's superior durability and cost-effectiveness, factors critical in resource-limited settings, their overriding selection of composite for its esthetic qualities and conservative cavity preparation warrants critical exploration.

This student preference is not purely a reflection of material performance but is likely shaped by powerful institutional and cultural drivers. Modern undergraduate curricula, influenced by global trends and significant patient demand for metal-free restorations, often place a heavier emphasis on composite protocols.¹² This pedagogical focus likely leads to greater student proficiency and confidence with resin materials, which subsequently translates into a clinical preference. This finding underscores a potential curricular bias that may not fully align with the contextual clinical needs of a resource-limited setting like Pakistan, where long-term affordability

and longevity under challenging conditions remain paramount.

These results contribute meaningfully to the global discourse on the phase-down of dental amalgam, aligning with the objectives of the Minamata Convention on Mercury, which encourages a reduction in amalgam use due to environmental and health concerns. The clinical decision between amalgam and composite has evolved into a more nuanced judgment, factoring not only material properties but also clinical indication, patient expectations, and health system capabilities. For instance, a meta-analysis published in the UK demonstrated that posterior composites performed comparably to amalgam in terms of survival rates over five years when placed with appropriate technique and isolation protocols.¹³ Similarly, research from Brazil and Nigeria, both middle-income countries, has shown that composite resins, under optimal conditions, can achieve restoration longevity comparable to amalgam, particularly when dentists receive adequate training and access to high-quality materials.¹⁴ Nevertheless, in low-resource environments, the technique sensitivity of composites remains a significant barrier. They require strict moisture control, incremental placement, and high-quality curing lights, which may not be consistently available in public-sector dental clinics. Studies found that composite restorations had higher failure rates in public health settings due to inadequate operator training and limited infrastructure.¹⁵ In such contexts, the continued use of dental amalgam, valued for its durability, ease of placement, and minimal equipment dependency, may still be a pragmatic choice, especially for large Class I and II restorations in posterior teeth.

Conversely, a critical perspective requires acknowledging the distinct clinical contexts where amalgam retains superior performance and practical advantage. Numerous clinical trials and



retrospective studies consistently highlight amalgam's reliability in high-stress conditions, such as large multi-surface Class II restorations, cases involving bruxism, and treatment of high caries-risk patients. A longitudinal study, which followed 1,748 restorations over seven years, found that amalgam restorations had significantly lower failure rates than composites in posterior teeth, particularly in complex multi-surface applications.¹⁵ Likewise, the UK-based Dental Practice Board survey observed that amalgam restorations exhibited longer survival times than composite counterparts, especially in patients with suboptimal oral hygiene.¹³ In moisture-compromised settings, common in pediatric or geriatric care, or in facilities without rubber dam isolation, amalgam's tolerance to minor contamination offers a major procedural advantage. This is particularly pertinent in resource-constrained health systems like Pakistan, where infrastructural limitations may restrict consistent access to high-quality curing lights, matrix systems, or trained auxiliary staff. A recent regional analysis emphasized that the cost-effectiveness and low technique sensitivity of amalgam remain vital in South Asian public dental clinics, where patient volume is high and operator time is limited. In this context, the affordability and proven longevity of amalgam, often exceeding 10–15 years with minimal maintenance, position it as an indispensable material.⁸ Therefore, curriculum design must reflect this dual reality: promoting the transition to modern, esthetic materials while preserving competence in the use of amalgam where clinically justified.

Another important dimension in the amalgam versus composite debate relates to the presence of mercury in dental amalgam. Although amalgam has been considered safe for over a century of use, concerns regarding mercury release, occupational exposure for dental personnel, and potential environmental impact have gained prominence in recent years. While the actual health risk to patients from amalgam restorations remains minimal according to most scientific assessments, the environmental implications of mercury disposal and patient preference for metal-free, esthetic restorations have contributed significantly to the increasing adoption of composites.¹⁶ This trend reflects not only material performance but also broader public health and environmental considerations that continue to shape restorative practices worldwide.

Given the findings regarding student preference and the practical realities of resource-

constrained dentistry, we offer strong recommendations for curriculum development within Pakistani dental institutions: (i) Contextualized Material Training: While composite training must be modernized and emphasized, the curriculum must maintain robust, evidence-based training in Amalgam hygiene and placement for large, demanding, or high caries-risk restorations, acknowledging its contextual necessity. (ii) Critical Thinking in Selection: Educational strategies should focus on fostering critical thinking in material selection, teaching students to perform a rapid risk-benefit analysis that incorporates factors unique to resource-constrained settings (e.g., patient adherence, long-term maintenance costs, and moisture control difficulty) (iii) Protocol-Driven Clinical Pathways: Institutions should clearly delineate protocol-driven clinical pathways to guide students in selecting the appropriate material based on clinical indication rather than aesthetic preference alone, ensuring graduates are equipped to deliver both esthetic and durable care tailored to diverse socioeconomic realities.

In addition to the inherent limitations of using self-report measures (susceptible to social desirability and recall bias), the scope of this study's generalizability is restricted by the sampling strategy employed. The use of a convenience sample drawn exclusively from a single institutional setting introduces potential sampling bias. Therefore, caution must be exercised when extrapolating these findings beyond the specific context of this study. To enhance the robustness and external validity of future investigations, a multi-center study incorporating random or stratified sampling across geographically and institutionally diverse sites is strongly recommended. Such an approach would mitigate institution-specific biases and provide a more comprehensive, generalizable understanding of the phenomena under investigation. The robust findings of this study are further supported by the internal consistency demonstrated by the questionnaire/scale in our sample (Cronbach's alpha = 0.82), confirming its credibility and fitness for purpose within this specific population.

A further limitation of the current study is the lack of data regarding the participants' clinical experience with the materials assessed. While we measured student preference and perception, the manuscript does not quantify the actual volume of amalgam versus composite restorations that students had placed before completing the survey. Without incorporating these objective clinical exposure metrics, it is challenging to definitively

contextualize whether the observed preferences are driven by theoretical knowledge, curriculum emphasis, or genuine hands-on experience and confidence. Future research should incorporate detailed clinical logs or portfolios to correlate material preferences directly with the depth and breadth of practical exposure.

Conclusion

This multi-provincial study shows a clear preference among Pakistani dental students for composite resin over amalgam for posterior restorations, mainly due to esthetics, conservative preparation, and curricular emphasis, despite recognizing amalgam's durability and cost-effectiveness. The findings highlight the need for an evidence-based, balanced curricular approach that equips future dentists to select materials based on clinical indication, patient needs, and long-term outcomes rather than preference alone.

Acknowledgments

The authors thank the administration and faculty of Islamic International Dental College (IIDC), Riphah International University, for their support during this study. Special appreciation is extended to the students whose participation and valuable insights made this research possible.

Author Contribution

UH and AA conceived the study and designed the work. AA, MA, and SS collected the data. MA, RS, and MS analyzed and interpreted the data and generated themes. All the authors contributed to the drafting and writing of the manuscript. UH, AA, MA, SS, MS, and RS critically reviewed and revised the manuscript. All authors proofread the finalized manuscript.

Data Availability Statement

All relevant data are within the manuscript. Additional data supporting this study are available from the corresponding author upon reasonable request.

Funding

No external funding was received.

Conflict of Interest

None.

References

- Ghods S, Arzani S, Shekarian M, Aghamohseni M. Cement selection criteria for full coverage restorations: A comprehensive review of literature. *Journal of clinical and experimental dentistry*. 2021;13(11):e1154.
- Bailey OS. Amalgam phase-out: what next for dentistry? Costs and benefits of the alternative direct restorations: Newcastle University; 2025.
- Mackenzie L. Dental amalgam: a practical guide. *Dental Update*. 2021;48(8):607-18.
- Klaisiri A, Phumpatrakom P, Thamrongananskul N. Chemical surface modification methods of resin composite repaired with resin-modified glass-ionomer cement. *European Journal of Dentistry*. 2023;17(03):804-8.
- Mittal N, Gupta S. Compomers: A Review of Literature. *Acta Scientific Dental Sciences (ISSN: 2581-4893)*. 2021;5(6).
- Paolone G, Baldani S, De Masi N, Mandurino M, Collivasone G, Scotti N, et al. Translucency of bulk-fill composite materials: A systematic review. *Journal of Esthetic and Restorative Dentistry*. 2024;36(7):995-1009.
- Hotchandani D, Sajjanar D, Lakshmi Y, Peri V, Banik A, Duggal S. Cention-N: a comprehensive review. *Eur Chem Bull*. 2023;12:5528-36.
- Fatima F, Shah SAU, Ameer S, Alim S, Malik S, Khaliq S. Comparison of Clinical Outcome of Direct Composite Resin Versus Amalgam Restoration in Permanent Posterior Teeth. *Journal of Health and Rehabilitation Research*. 2024;4(2):1-4.
- Pawar LP, Patel A, Chandak M, Wazurkar S, Nadgouda M. Perfecting the Craft: Composite Restoration Elevated With the Stamp Technique. *Cureus*. 2024;16(9).
- Zehra A, Rafique MU, Sohaib M, Ullah H, Khan S, Hussain M. Effectiveness of Resin-Composite Fillings vs. Amalgam Fillings in Restorative Dentistry: A Meta-Analysis on Durability and Patient Satisfaction. *Indus Journal of Bioscience Research*. 2025;3(3):63-8.
- Pani SC, Al Abbassi MF, Al Saffan AD, Al Sumait MA, Shakir AN. Factors influencing Saudi dental students' preference of amalgam or composite for posterior dental restorations. *Saudi Journal of Oral Sciences*. 2014;1(1):30-6.
- Daood U, Sidhu P, Ab Malik N, Lin SL, Wilson NH, Lynch CD, et al. Current and future trends in the teaching of direct posterior composites restorations at dental schools in Canada: a cross-sectional study. *Journal of Dentistry*. 2025;105818.
- Santos M, Saadaldin S, Rêgo H, Tayan N, Melo N, El-Najar M. Clinical Longevity of Complex Direct Posterior Resin Composite and Amalgam Restorations: A Systematic Review and Meta-analysis. *Operative Dentistry*. 2025.
- Kisumbi BK, Osiro OA, Gathece LW, Maina SW. Dental Amalgam Phase-Down—Status, Alternatives, Strategies and Preparedness for Implementation: A Review. *International Journal of Dentistry*. 2025;2025(1):6688410.
- Moraes RR, Cenci MS, Moura JR, Demarco FF, Loomans B, Opdam N. Clinical performance of resin composite restorations. *Current Oral Health Reports*. 2022;9(2):22-31.
- Geier D, Geier M. Dental amalgam fillings and mercury vapor safety limits in American adults. *Human & Experimental Toxicology*. 2022;41:09603271221106341.